

v

AD-A179 713

OFFICE OF NAVAL RESEARCH

Contract N00014-K-0043

R & T Code 413f0001---01

TECHNICAL REPORT No. 35

Semiempirical Study of Polyatomic Rare Gas Halides: Application to the Xe_nCl Systems

by

Isidore Last and Thomas F. George

Prepared for Publication

in

Journal of Chemical Physics Letters

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

April 1987

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

DTIC
ELECTE
APR 27 1987
S D
E

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS			
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited			
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE						
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/87/TR-35			5. MONITORING ORGANIZATION REPORT NUMBER(S)			
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NOS.			
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT NO.
11. TITLE Semiempirical Study of Polyatomic Rare Gas Halides: Application to the Xe _n Cl Systems						
12. PERSONAL AUTHOR(S) Isidore Last and Thomas F. George						
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) March 1987		15. PAGE COUNT 40
16. SUPPLEMENTARY NOTATION Prepared for Publication in Chemical Physics Letters						
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)			
FIELD	GROUP	SUB. GR.	POLYATOMIC, RARE GAS HALIDES, Xe₂Cl, Xe₃Cl and Xe₁₂Cl Xe-Cl SYSTEMS , Xenon Cl_n Cl_n DIATOMICS, DIATOMIC SYSTEMS , QUASISTABLE, Xe ₂ Cl ⁺ , POLARIZATION FORCES			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) In order to calculate different electronic states in polyatomic rare gas-halogen systems, including ionic states with charge transfer, a semiempirical diatomics-in-ionic systems (DIIS) method is presented. In this method the Hamiltonian matrix elements are expressed in terms of diatomic potentials, except for those which are responsible for coupling between the ionic and neutral states. The coupling matrix elements are determined by fitting the diatomic transition moments. Calculations are performed for Xe ₂ Cl, Xe ₃ Cl and Xe ₁₂ Cl. The results reveal at least three quasistable ionic Xe ₂ Cl ⁺ molecules. Such molecules can attract more Xe atoms, which are binded mainly by polarization forces.						
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input checked="" type="checkbox"/> DTIC USERS <input type="checkbox"/>				21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE NUMBER (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL	

Semiempirical study of polyatomic rare gas
halides: Application to the Xe_nCl systems

Isidore Last* and Thomas F. George
Department of Physics & Astronomy
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or Special
A-1	

Abstract

In order to calculate different electronic states in polyatomic rare gas-halogen systems, including ionic states with charge transfer, a semiempirical diatomics-in-ionic-systems (DIIS) method is presented. In this method the Hamiltonian matrix elements are expressed in terms of diatomic potentials, except for those which are responsible for coupling between the ionic and neutral states. The coupling matrix elements are determined by fitting the diatomic transition moments. Calculations are performed for Xe₂Cl, Xe₃Cl and Xe₁₂Cl. The results reveal at least three quasistable ionic Xe₂⁺Cl⁻ molecules. Such molecules can attract more Xe atoms, which are binded mainly by polarization forces.



*Permanent address: Soreq Nuclear Research Center, Yavne 70600, Israel

I. Introduction

Interest in rare gas-halogen systems has been increasing due to their spectroscopic, and in particular lasing, properties.¹⁻¹² The lasing properties are associated with the formation of quasistable ionic molecules R^+X^- and $R_2^+X^-$, where R and X are the rare gas and halogen atoms, respectively. These ionic molecules are obtained in rare gas-halogen gas mixtures either by electron impact^{1-4,7,8} or by optical, most often laser, irradiation.^{5,9,11,12} Recently the ionic rare gas-halogen molecules $Xe_2^+Cl^-$ were detected also in rare gas solids doped by Cl_2 and HCl.^{13,14} Rare gas-halogen systems are of interest not only for the configurations where the quasistable ionic R^+X^- and $R_2^+X^-$ molecules are formed, but also for configurations of van der Waals RX , RX_2 or RXH complexes.¹⁵⁻¹⁹

The theoretical consideration of the rare gas-halogen systems involves a knowledge of potential curves and potential energy surfaces (PES). The calculation of these PES is the aim of the present paper. For the sake of generalization, we shall consider here arbitrary R_nX systems where n may be larger than 2 and consequently the number of atoms larger than 3. Although the RX and R_2X systems have been mostly the subject of study, the R_nX systems with $n > 2$ can also be of interest due to the possibility of the formation of cluster-like molecules or large van der Waals complexes. The $R_n^+X^-$ ($n > 2$) excited systems can also be formed in rare gas solids doped by halogen molecules.²⁰

In the case of diatomic RX ($n = 1$) systems some of the potential curves for different electronic states are well known from ab initio calculations.²¹⁻²³ The RX potential curves, mostly for the ground state, were studied also by applying model approach²⁴ and by using experimental data.^{4,15,26} The information concerning the triatomic R_2X systems, not to

mention the R_nX systems for $n > 2$, is much more scarce. The experimental data of the R_2X systems are limited mainly to the transition energies.^{7,9,11,12} The all-electron ab initio calculations were performed for the Ar_2F and Kr_2F systems only.²⁷ The ab initio calculation of the Xe_2Cl system which contains many more electrons was performed by using a simplified model.²⁸ The R_2X systems were also calculated semiempirically by using the diatomics-in-molecules (DIM) method.^{29,30}

The advantages of the semiempirical approach in the PES calculations of polyatomic systems containing a large number of electrons are well known. In particular, the DIM method is often used in the PES calculations.³⁷ Unfortunately, in the case where ionic states are involved, the application of the DIM method becomes much more complicated if the coupling between the neutral and ionic states is taken into account.³² The difficulties arise largely due to the "zero overlap of atomic orbitals" approximation which is usually used in the DIM method. According to this approximation, the overlap between atomic orbitals belonging to different atoms is neglected. Consequently the off-diagonal matrix elements between any electronic configurations with different numbers of electrons on the same atom, like the neutral and ionic states, are equal to zero, such that the states are uncoupled. This shortcoming of the usual DIM method can be overcome by introducing additional assumptions and new parameters.³²⁻³⁴ However, such an approach increases the complexity of the calculation and makes the results more arbitrary. In DIM calculations of the R_2X systems,^{29,30} the usual version of the method was used with the off-diagonal matrix elements between neutral and ionic states equal to zero and the independent consideration of the R_2X and R_2X^+ states.

The influence of the coupling between the neutral and ionic states of the $R_n X$ systems is not quite clear. This problem was studied only in the connection with ab initio calculations of the XeX diatomic systems.²² According to this study, the coupling is of the order of 10% for the equilibrium XeX van der Waals distance and decreases slowly as the interatomic distance increases. It is difficult to estimate the influence of the coupling on different properties of the RX systems, although it obviously reduces size of the van der Waals RX molecule. For example, the van der Waals radii of the Xe and Cl atoms are 2.2 Å and 7.8 Å,³⁵ respectively, and while the XeCl van der Waals distance is expected to be 4.0 Å, in reality it is only 3.23 Å.²⁶ Because the coupling may be different for different rare gas-halogen systems, the use of diatomic RX potential curves in the semiempirical calculations of polyatomic systems (DIM, for example) may lead to confusing results. Thus, the coupling and consequently the interaction are expected to be small in the case of a RX_2 van der Waals complex due, partly, to lower electron affinity of the X_2 molecule compared to the X atom,²⁰ which is confirmed by the spectroscopic properties of X_2 molecules trapped in a rare gas matrix.³⁶ However, neglecting the change in the coupling and using the empirical RX interaction in the DIM calculation of the van der Waals RX_2 molecule, one would obtain the RX_2 interaction as strong, roughly, as in the R-X case, in contradiction with the experiment. The coupling between neutral and ionic states must be taken into account if one wants to calculate the off-diagonal (transition) dipole moments.

We shall present in this paper a semiempirical method which is designed for a calculation of rare gas-halogen systems $R_n X$ consisting of an arbitrary number of rare gas atoms and one halogen atom. Introducing some

modifications, the method can be extended to R_nHX systems. The method takes into account in a straightforward way the coupling between the neutral R_nX and the ionic $R_n^+X^-$ configurations, as well as the positive charge delocalization in the ionic $R_n^+X^-$ states. The method makes use of diatomic potentials as initial data for a calculation, but in a different way than the DIM method. A main feature of the method is the consideration of the systems with ionic states, and hence we call it the "diatomics-in-ionic-systems" (DIIS) method.

The spin-orbit coupling in the Cl atom and Xe^+ ion will be not taken into account directly. The shift of energy levels resulting from this coupling can always be incorporated indirectly by using proper empirical or ab initio potentials. The main disadvantage of the neglect of spin-orbit coupling is the loss of some of the excited states. The direct inclusion of spin-orbit coupling does not present theoretical obstacles. However, it would make the Hamiltonian matrix elements much more complicated and would increase significantly the order of the matrix. The latter is of primary concern for large systems, like rare gas solids with halogens.²⁰

The theoretical consideration of the DIIS method is given in Section II. Section III presents the equations for the diatomic terms which determine the matrix elements of the method. The diatomic terms for the Xe_nCl systems are given in Section IV. Section V deals with the calculation of the Xe_nCl systems, namely Xe_2Cl , Xe_3Cl and $Xe_{12}Cl$, and Section VI is the Conclusion.

II. Diatomics-in-ionic-systems method

A polyatomic rare gas-halogen system R_nX will be presented here as $R_1R_2\dots R_{J-1}X_J$ ($J = n + 1$), where the R_i atoms have a closed shell in the neutral state and P symmetry shell in the ionized state R_i^+ , whereas the X_J

atom has a P symmetry shell in the neutral state and a closed shell when becoming the negative ion X_J^- . The R_i atoms need not be just the rare gas atoms but also H atoms, with the difference that in the ionized state H^+ they are also spherically symmetric. The systems with H atoms can be treated as a particular case which we shall not consider separately, suggesting that all R_i^+ ions have a P symmetry shell.

The wave function of the polyatomic system in a fixed electronic configuration will be presented here as an antisymmetrized product of atomic group functions,³⁷ which are assumed to be exact many-electronic wave functions of individual atoms,

$$\phi_{m_1 \dots m_J} = \hat{A} \prod_i^J \chi_{im_i}, \quad (1)$$

where \hat{A} is the operator which antisymmetrized electrons belonging to different group functions χ_{im_i} and m_i are the atomic state indexes. The atomic group functions in the polyatomic wave function (1) are not orthogonal one to another which makes the energy expression very complicated.^{38,39} The consideration is significantly simplified by using the zero overlap of atomic orbitals (ZAO) approximation which is very common in semiempirical methods, in particular in the DIM method.³¹ The ZAO approximation allows one to omit the antisymmetrization operator \hat{A} , thus rendering the polyatomic wave function as a simple product of atomic group functions. The mistake arising from the neglect of the overlap is proportional to the square of the interatomic overlap integrals³⁹ and has been shown to be small, at least in DIM calculations.⁴⁰

In the accepted approximations, the wave function of a neutral $R_1 R_2 \dots R_{J-1} X_J$ electronic configuration is equal to the product of S

symmetrical (closed shell) $\chi_i^{(0)}$ group functions of the R_i atoms and the P symmetrical $\chi_{Jm}^{(0)}$ group function of the X_J (halogen) atom,

$$\phi_{Jm} = \prod_i^{J-1} \chi_i^{(0)} \chi_{Jm}^{(0)}, \quad m = 1, 2, 3 \quad (2)$$

where the index m indicates the orientation of the $\chi_{Jm}^{(0)}$ group function along the x, y, z -axes. In an ionic state, one of the $J-1$ rare gas atoms (R_i^+) has a positive charge and is distinguished by its $\chi_{im}^{(+)}$ group function orientation, whereas the $\chi_J^{(-)}$ group function of the halogen ion X_J^- has S symmetry,

$$\phi_{im} = \prod_j^{J-1} \chi_j^{(0)} \chi_{im}^{(+)} \chi_J^{(-)}, \quad i=1, 2, \dots, J-1; m=1, 2, 3; j \neq i. \quad (3)$$

The upper indexes (0), (+) and (-) indicate in Eqs. (2)-(3) and below the charge of an atom. It is important to emphasize that Eqs. (1)-(3) do not necessarily imply the one-electron approximation, since the group functions are arbitrary and can include correlation between electrons belonging to the same atom. Using the group functions product, we neglect only the interatomic correlation which describes, in particular, the van der Waals attraction. However, in the framework of the semiempirical approach, the van der Waals interaction can be taken into account indirectly by using empirical diatomic potentials.

The three functions [Eq.(2)] of the neutral configurations and the $3(J-1)$ functions [Eq.(3)] of the ionic configurations form the polyatomic basis set. The linear combination of these functions forms the wave function of the system

$$\phi = \sum_{i=1}^J \sum_{m=1}^3 c_{im} \phi_{im}. \quad (4)$$

The energy levels of this wave function are obtained now in the usual way by forming the Hamiltonian matrix of order $3J \times 3J$, where J is the number of atoms in the system.

Due to the lack of interatomic electron permutations in the polyatomic functions of Eqs.(2) and (3), the Hamiltonian of the system can be partitioned into atomic and interatomic terms. This partitioning depends on the representation of the polyatomic wave function and changes from one matrix element to another. The Hamiltonian which is used for the evolution of the matrix elements for the neutral states has the form

$$\hat{H} = \sum_j^{J-1} \hat{h}_j(v_j) + \sum_{j_1}^{J-2} \sum_{j_2(>j_1)}^{J-1} h_{j_1 j_2}(v_{j_1} + v_{j_2}) + \hat{h}_J(v_J) + \sum_j^{J-1} h_{jJ}(v_j + v_J), \quad (5)$$

where the \hat{h}_j and \hat{h}_J terms with one index include all energy operators of the electrons belonging to one atom, the $h_{j_1 j_2}$ and h_{jJ} terms with two indexes include the interactions between electrons and nuclei belonging to different atoms, and v_j indicates the number of electrons in atom j . We will present the energy of the system relative to the energy of separated on infinity atoms

$$E_\infty = \sum_j^{J-1} E_j^{(0)} + E_J^{(0)}, \quad (6)$$

where $E_j^{(0)}$ and $E_J^{(0)}$ are the energies of the rare gas and halogen atoms, respectively. Using the expressions (2), (5) and (6), one obtains the following expression for the diagonal matrix elements of the neutral states:

$$H_{J_m, J_m} = \int dv \phi_{J_m} \hat{H} \phi_{J_m} = \sum_{j_1}^{J-2} \sum_{j_2(>j_1)}^{J-1} U_{j_1, j_2}^{(00)} + \sum_j^{J-1} U_{j, J_m}^{(00)}, \quad m=1, 2, 3. \quad (7)$$

In Eq. (7), $U_{j_1, j_2}^{(00)}$ and $U_{j, Jm}^{(00)}$ are the potentials of the interaction between two rare gas atoms and between a rare gas atom and a halogen atom, respectively:

$$U_{j_1, j_2}^{(00)} = \int dv [x_{j_1}^{(0)}]^2 h_{j_1 j_2} [x_{j_2}^{(0)}]^2, \quad (8)$$

$$U_{j, Jm}^{(00)} = \int dv [x_j^{(0)}]^2 h_{jJ} [x_{Jm}^{(0)}]^2. \quad (8')$$

In the expression for the off-diagonal matrix elements of the neutral states, all integrals with one-atom operators \hat{h}_j and \hat{h}_J and with the interatomic terms $h_{j_1 j_2}$ vanish either due to the difference in the symmetry of functions with different m or due to the ZAO approximation. Only the h_{jJ} terms give a non-zero contribution,

$$H_{Jm_1, Jm_2} = \int dv \phi_{Jm_1} \hat{H} \phi_{Jm_2} = \sum_j^{J-1} \int dv [x_j^{(0)}]^2 h_{jJ} x_{Jm_1}^{(0)} x_{Jm_2}^{(0)}. \quad (9)$$

Let us consider now the diagonal matrix elements for the ionic states [Eq.(3)]. The Hamiltonian for the ϕ_{im} function which describes the state with a positive charge on the i -th rare gas atom and a negative charge on the J -th (halogen) atom has the form

$$\hat{H} = \sum_j^{J-1} \hat{h}_j(v_j) + \sum_{j_1}^{J-2} \sum_{j_2(>j_1)}^{J-1} h_{j_1 j_2}(v_{j_1} + v_{j_2}) + \hat{h}_i(v_i - 1) + \sum_j^{J-1} h_{ij}(v_i - 1 + v_j) + \hat{h}_J(v_J + 1) + \sum_j^{J-1} h_{jJ}(v_j + v_J + 1) + h_{iJ}(v_i + v_J), \quad j, j_1, j_2 \neq i. \quad (10)$$

This form includes terms which are responsible for the polarization of neutral atoms by the ions R_i^+ and X_J^- . It is more convenient to determine the energy of polarization in the electrostatic approximation, separating it as

a special term ϵ_i ,⁴¹

$$H_{im,im} = \bar{H}_{im,im} + \epsilon_i. \quad (11)$$

Considering atoms and ions as non-overlapping spheres, one obtains

$$\epsilon_i = - \sum_j^{J-1} C_j \left(\frac{\vec{R}_{ij}}{R_{ij}^3} - \frac{\vec{R}_{ji}}{R_{ji}^3} \right)^2 + \Delta_i, \quad j \neq i, \quad (12)$$

$$C_j = e^2 \alpha_j / 2, \quad (13)$$

where α_j are the rare gas atom polarizabilities, \vec{R}_{ij} and \vec{R}_{ji} are the vectors between the corresponding atoms, and Δ_i includes the dipole-dipole interaction. The matrix elements $\bar{H}_{im,im}$ are obtained by using the Hamiltonian without polarization terms $-C_j/R_{ij}^4$:

$$\begin{aligned} \hat{H} = & \sum_j^{J-1} \bar{h}_j(v_j) + \sum_{j_1}^{J-2} \sum_{j_2(>j_1)}^{J-1} h_{j_1 j_2}(v_{j_1} + v_{j_2}) + \bar{h}_i(v_i - 1) + \sum_j^{J-1} \bar{h}_{ij}(v_i - 1 + v_j) + \\ & \bar{h}_j(v_j + 1) + \sum_j^{J-1} \bar{h}_{jJ}(v_j + v_J + 1) + h_{iJ}(v_i + v_J), \end{aligned} \quad (10')$$

$$\bar{h}_{ij} = h_{ij} - (-C_j/R_{ij}^4) \quad (14)$$

Using the expressions (3), (6) and (10'), one obtains

$$\begin{aligned} \bar{H}_{im,im} = & \int dv \phi_{im} \hat{H} \phi_{im} = I_i - A_J + \sum_{j_1}^{J-2} \sum_{j_2(>j_1)}^{J-1} U_{j_1 j_2}^{(00)} + \\ & \sum_j^{J-1} (U_{im,j}^{(+0)} + U_{J,j}^{(-0)}) + U_{im,J}^{(+-)}, \quad j_1 j_2 \neq i, \end{aligned} \quad (15)$$

$$U_{im,j}^{(+0)} = \int dv [\chi_{im}^{(+)}]^2 \bar{h}_{ij} [\chi_j^{(0)}]^2 = U_{im,j}^{(+0)} + C_j/R_{ij}^4, \quad i=1,2..J-1, j \neq i. \quad (16)$$

The off-diagonal matrix elements between the ionic states with the same localization of the positive charge but different orientation m are obtained

in the same way as the matrix elements in Eq.(9),

$$H_{im_1, im_2} = \int dv \phi_{im_1} \hat{H} \phi_{im_2} = \sum_j^{J-1} \int dv [\chi_j^{(0)}]^2 \bar{h}_{ij} \chi_{im_1}^{(+)} \chi_{im_2}^{(+)} + \int dv [\chi_j^{(-)}]^2 \bar{h}_{ij} \chi_{im_1}^{(+)} \chi_{im_2}^{(+)}, \quad j \neq i. \quad (17)$$

The off-diagonal matrix elements between the ionic states with different localization of the positive charge and between the neutral and ionic states are also obtained in a similar way,

$$H_{i_1 m_1, i_2 m_2} = \int dv \chi_{i_1}^{(0)} \chi_{i_2 m_2}^{(+)} \hat{H}_{i_1 i_2} (v_{i_1} + v_{i_2} - 1) \chi_{i_1 m_1}^{(+)} \chi_{i_2}^{(0)}, \quad (18)$$

$$H_{J m_1, im_2} = \int dv \chi_{J m_1}^{(0)} \chi_i^{(0)} \hat{H}_{J i} (v_i + v_J) \chi_J^{(-)} \chi_{im_2}^{(+)}, \quad (19)$$

where $\hat{H}_{i_1 i_2} (v_{i_1} + v_{i_2} - 1)$ is the Hamiltonian of the charged $(R_{i_1} R_{i_2})^+$ diatomic fragment and $\hat{H}_{J i} (v_i + v_J)$ is the Hamiltonian of the $R_i X_J$ diatomic fragment. The exchange integral (18) presents the valence interaction between two rare gas atoms with a common +1 charge. The exchange integral (19) is responsible for the coupling between the neutral and ionic states of the rare gas-halogen diatomic systems.

The matrix elements obtained above are expressed by diatomic terms which are determined by atomic group functions oriented, when of P symmetry, along Cartesian (x,y,z) axes. In order to express the matrix elements by the diatomic terms of usual Σ and Π orientations, one needs to transform the P symmetrical group functions to a new set of functions

$$\chi_{im} = \sum_{k=1}^3 S_{ij, mk} \psi_{ik, j}, \quad (20)$$

where the indices $m=1,2,3$ indicate the x-, y- and z-orientations, respectively, $\psi_{i1, j}$ is the Σ group function of the i-th atom oriented along

the \vec{R}_{ij} vector connecting the two atoms, and $\psi_{12,j}$ and $\psi_{13,j}$ are the group functions of the i -th atom oriented perpendicular to the \vec{R}_{ij} -vector. We will choose one of these two perpendicular orientations (Π) as located in the plane formed by the \vec{R}_{ij} -vector and the x -axis. The second orientation which is perpendicular to this plane will be denoted by $\bar{\Pi}$. The 3×3 $S_{ij,mk}$ -matrix has the form

$$\begin{aligned} S_{ij,11} &\equiv S_{ij,x\Sigma} = (x_j - x_i)/R_{ij}, & S_{ij,12} &\equiv S_{ij,x\Pi} = r_{ij}/R_{ij}, \\ S_{ij,13} &\equiv S_{ij,x\bar{\Pi}} = 0 \end{aligned} \quad (21)$$

$$\begin{aligned} S_{ij,21} &\equiv S_{ij,y\Sigma} = (y_j - y_i)/R_{ij}, & S_{ij,22} &\equiv S_{ij,y\Pi} = \\ &-(x_j - x_i)(y_j - y_i)/(R_{ij}r_{ij}), & S_{ij,23} &\equiv S_{ij,y\bar{\Pi}} = (z_j - z_i)/r_{ij} \end{aligned} \quad (21')$$

$$\begin{aligned} S_{ij,31} &\equiv S_{ij,z\Sigma} = (z_j - z_i)/R_{ij}, & S_{ij,32} &\equiv S_{ij,z\Pi} = \\ &-(x_j - x_i)(z_j - z_i)/(R_{ij}r_{ij}), & S_{ij,33} &\equiv S_{ij,z\bar{\Pi}} = -(y_j - y_i)/r_{ij}, \end{aligned} \quad (21'')$$

where R_{ij} is the interatomic distance and r_{ij} is the projection of the R_{ij} vector on the yz -plane,

$$R_{ij} = \sqrt{(x_j - x_i)^2 + (y_j - y_i)^2 + (z_j - z_i)^2} \quad (22)$$

$$r_{ij} = \sqrt{(y_j - y_i)^2 + (z_j - z_i)^2}. \quad (22')$$

Using the transformation (21) we will express the matrix elements (7), (9), (15), (17), (18) and (19) by Σ and Π diatomic terms,

$$\begin{aligned} H_{Jm,Jm} &= \sum_{j_1}^{J-2} \sum_{j_2(>j_1)}^{J-1} U_{j_1,j_2}^{(00)} + \sum_j^{J-1} [S_{Jj,m\Sigma}^2 U_{Jj,\Sigma}^{(00)} \\ &+ (S_{Jj,m\Pi}^2 + S_{Jj,m\bar{\Pi}}^2) U_{Jj,\Pi}^{(00)}] \end{aligned} \quad (23)$$

$$\begin{aligned}
H_{im, im} = & I_i - A_J + \sum_{j_1}^{J-2} \sum_{j_2(>j_1)}^{J-1} U_{j_1, j_2}^{(00)} + \sum_j^{J-1} [S_{ij, m\Sigma}^2 U_{ij, \Sigma}^{(+0)} \\
& + (S_{ij, m\Pi}^2 + S_{ij, m\bar{\Pi}}^2) U_{ij, \Pi}^{(+0)}] + S_{iJ, m\Sigma}^2 U_{iJ, \Sigma}^{(+0)} \\
& + (S_{iJ, m\Pi}^2 + S_{iJ, m\bar{\Pi}}^2) U_{iJ, \Pi}^{(+0)} + \sum_j^{J-1} U_{jJ}^{(0-)} + \epsilon_i, \quad j, j_1, j_2 \neq i \quad (24)
\end{aligned}$$

$$\begin{aligned}
H_{Jm_1, Jm_2} = & \sum_j^{J-1} [S_{Jj, m_1 \Sigma} S_{Jj, m_2 \Sigma} U_{Jj, \Sigma}^{(00)} \\
& + (S_{Jj, m_1 \Pi} S_{Jj, m_2 \Pi} + S_{Jj, m_1 \bar{\Pi}} S_{Jj, m_2 \bar{\Pi}}) U_{Jj, \Pi}^{(00)}], \quad (25)
\end{aligned}$$

$$\begin{aligned}
H_{im_1, im_2} = & \sum_j^{J-1} [S_{ij, m_1 \Sigma} S_{ij, m_2 \Sigma} U_{ij, \Sigma}^{(+0)} + (S_{ij, m_1 \Pi} S_{ij, m_2 \Pi} \\
& + S_{ij, m_1 \bar{\Pi}} S_{ij, m_2 \bar{\Pi}}) U_{ij, \Pi}^{(+0)}] + S_{iJ, m_1 \Sigma} S_{iJ, m_2 \Sigma} U_{iJ, \Sigma}^{(+0)} + (S_{iJ, m_1 \Pi} S_{iJ, m_2 \Pi} \\
& + S_{iJ, m_1 \bar{\Pi}} S_{iJ, m_2 \bar{\Pi}}) U_{iJ, \Pi}^{(+0)}, \quad j \neq i \quad (26)
\end{aligned}$$

$$\begin{aligned}
H_{i_1 m_1, i_2 m_2} = & S_{i_1 i_2, m_1 \Sigma} S_{i_2 i_1, m_2 \Sigma} V_{i_1 i_2, \Sigma} + (S_{i_1 i_2, m_1 \Pi} S_{i_2 i_1, m_2 \Pi} + \\
& S_{i_1 i_2, m_1 \bar{\Pi}} S_{i_2 i_1, m_2 \bar{\Pi}}) V_{i_1 i_2, \Pi} \quad (27)
\end{aligned}$$

$$\begin{aligned}
H_{Jm_1, im_2} = & S_{Ji, m_1 \Sigma} S_{iJ, m_2 \Sigma} V_{Ji, \Sigma} + (S_{Ji, m_1 \Pi} S_{iJ, m_2 \Pi} \\
& + S_{Ji, m_1 \bar{\Pi}} S_{iJ, m_2 \bar{\Pi}}) V_{Ji, \Pi}. \quad (28)
\end{aligned}$$

The diatomic terms $V_{i_1 i_2}$ and V_{Ji} in the off-diagonal matrix elements (27) and (28) are expressed by the integrals (18) and (19), respectively, with the only difference that instead of x, y, z -oriented functions χ , the Σ, Π -oriented functions ψ of Eq. (20) are used. The diatomic U terms in the matrix elements (23)-(26) will be called Coulombic or diabatic potentials as they do not include exchange interactions presented by the off-diagonal

matrix elements (27)-(28). We have to note that the Coulombic terms U can include in a semiempirical way the van der Waals interaction which is not of Coulombic character.

The $3J \times 3J$ matrix H provides $3J$ eigenvalues E_k and $3J$ eigenvectors $c_{im}^{(k)}$ of the wave function of Eq.(4). Using the eigenvectors $c_{im}^{(k)}$, it is possible to calculate both static and transition moments. In the ZOA0 approximation the static dipole moment of any state is completely determined by the contribution of the ionic configuration. The x-component of the dipole moment $\vec{\mu}_k$ is

$$(\vec{\mu}_k)_x = e \sum_{i=1}^{J-1} w_i^{(k)} (\vec{R}_J - \vec{R}_i)_x \quad (29)$$

$$w_i^{(k)} = \sum_m (c_{im}^{(k)})^2, \quad (30)$$

where e is the electron charge and \vec{R}_J and \vec{R}_i are the radius vectors of the ions. The transition moment between a state with largely ionic configuration and a state with largely neutral configuration is

$$(\vec{\mu}_{kl})_x = e \sum_{i=1}^{J-1} w_i^{(kl)} (\vec{R}_J - \vec{R}_i)_x \quad (31)$$

$$w_i^{(kl)} = \sum_m c_{im}^{(k)} c_{im}^{(l)}. \quad (32)$$

III. Diatomic Coulombic and exchange terms

In the previous section the matrix elements of a polyatomic system were expressed by diatomic Coulombic (U) and exchange (V) terms. We shall now express these Coulombic and exchange terms by diatomic potentials which are assumed to be known. For the sake of simplicity, our study will be

restricted to systems with identical rare gas atoms R_i . There are four kinds of diatomic fragments: R_2 , R_2^+ , RX^- and RX . The expressions for the energy of any diatomic fragment will be obtained from the matrix elements of the previous section by removing all atoms to infinity except those which belong to this fragment.

A. R-R interaction

In the case of the $R_i R_j$ diatomic fragment, the ground state energy is equal to the diagonal matrix element [Eq.(23)] with one non-zero term only, namely $U_{ij}^{(00)}$. It follows that the diatomic term $U_{ij}^{(00)}$ is identical to the R_2 ground-state Σ potential

$$U_{ij}^{(00)} = \Sigma. \quad (33)$$

B. $(R-R)^+$ interaction

The energy of the diatomic fragment $(R_i R_j)^+$ is determined by a 2×2 matrix with non-zero off-diagonal matrix elements. In the case of the Σ -oriented R^+ wave function, the diagonal matrix element (24) and the off-diagonal matrix element (27) become

$$H_{11} = H_{22} = I_R - A_X + U_{ij,\Sigma}^{(+0)} - C_4/R_{ij}^4 = I_R - A_X + U_{ij,\Sigma}^{(+0)} \quad (34)$$

$$H_{12} = V_{ij,\Sigma}, \quad (34')$$

where $-C_4/R_{ij}^4$ is the polarization energy ϵ . The matrix elements (34) and (34') provide a ground-state potential Σ_g and an excited state potential Σ_u . Considering these potentials relative to the energy of separated ion R^+ and neutral atom R, one obtains

$$U_{ij,\Sigma}^{(+0)} = \frac{1}{2}(\Sigma_u + \Sigma_g), \quad U_{ij,\Sigma}^{(+0)} = U_{ij,\Sigma}^{(+0)} + C_4/R_{ij}^4 \quad (35)$$

$$V_{ij,\Sigma} = \frac{1}{2}(\Sigma_u - \Sigma_g). \quad (36)$$

Similar expressions are valid also for the Π states, with the only

difference that the lower state is π_u whereas the higher state is π_g ,

$$U_{ij,\pi}^{(+0)} = \frac{1}{2}(\pi_g + \pi_u), \quad U_{ij,\pi}^{(+0)} = U_{ij,\pi}^{(+0)} + C_4/R_{ij}^4 \quad (37)$$

$$V_{ij,\pi} = \frac{1}{2}(\pi_g - \pi_u). \quad (38)$$

C. R-X⁻ interaction

In the case of the $R_i X_j^-$ diatomic fragment the ground state energy is equal to the diagonal matrix element (24) with one non-zero diatomic term $U_{iJ}^{(0-)}$ along with the difference $I_R - A_X$ and the polarization energy term ϵ equal to $-C_4/R_{iJ}^4$. Denoting by Σ the RX^- ground state potential determined relatively to the separated R atom and X^- ion one obtains

$$U_{iJ}^{(0-)} = \Sigma, \quad U_{iJ}^{(0-)} = \Sigma + C_4/R_{iJ}^4 \quad (39)$$

D. R-X interaction

Because of the ionic-neutral coupling, the R-X diatomic potentials are not connected explicitly with the Coulomb terms U_{iJ} , like in the previous cases, but rather are obtained as the eigenvalues of a 2x2 matrix with the diagonal elements expressed by the Coulombic terms U_{iJ} and the off-diagonal elements equal to the exchange term V_{iJ} . In the case of the Σ -orientation, the matrix elements, derived from Eqs. (23),(24) and (28), are

$$H_{11} = U_{iJ,\Sigma}^{(00)} \quad (40)$$

$$H_{22} = I_R - A_X + U_{iJ,\Sigma}^{(+)} \quad (40')$$

$$H_{12} = V_{iJ,\Sigma} \quad (40'')$$

The matrix (40) allows us to express the Coulombic terms $U_{iJ,\Sigma}^{(00)}$ and $U_{iJ,\Sigma}^{(+-)}$ in terms of its eigenvalues, i.e., the RX ground-state (1Σ) and excited-state (2Σ) potentials which are assumed to be known:

$$U_{\Sigma}^{(00)} = \frac{1}{2}(1\Sigma + 2\Sigma - \beta) \quad (41)$$

$$U_{\Sigma}^{(+)} = \frac{1}{2}(1\Sigma + 2\Sigma + \beta) - (I_R - A_X) \quad (42)$$

$$\beta = \sqrt{(2\Sigma - 1\Sigma)^2 - 4V_{\Sigma}^2} \quad (43)$$

Similar expressions are valid for the Π -oriented potentials $U_{\Pi}^{(00)}$ and $U_{\Pi}^{(+)}$.

Equations (41)-(42) exclude from the R-X potentials the ionic-neutral coupling term V_{iJ} , which must be found independently. Before estimating this term, we shall consider the influence of the ionic-neutral coupling on the R-X properties. Due to this coupling, the ground state (1Σ) of the rare gas-halogen molecule is a mix of "pure" neutral and ionic states whose energies are determined by the diagonal matrix elements (40) and (40'). Because the ionization potential of the rare gas atom I_R is much higher than the electron affinity of the halogen atom A_X , the $H_{22} - H_{11}$ difference is large. It results in relatively weak coupling, at least for distances exceeding the van der Waals equilibrium distance.^{20,22} However, this coupling significantly affects some of the properties of the rare gas-halogen systems. These properties can be used to find the coupling terms by semiempirical fitting. In the present calculation the coupling term V_{iJ} is found by fitting the RX -to- R^+X^- transition moment which depends strongly on V_{iJ} and vanishes when V_{iJ} is zero.

IV. Diatomic terms of the Xe_nCl systems

For the Xe_nCl systems the diatomic fragments are: $Xe-Xe$, $(Xe-Xe)^+$, $Xe-Cl^-$ and $Xe-Cl$. The Coulombic potentials and the exchange terms of all of these diatomic fragments, except $Xe-Cl$, are expressed by diatomic Σ or Π potentials [see Eqs. (33)-(39)]. In the case of the $Xe-Cl$ diatomic fragment, as it was shown in the previous section, the Coulombic potentials U [Eqs.(41)-(42)] are expressed not only by diatomic potentials but also by the exchange (coupling) term V , which has to be found independently.

The diatomic potentials can be taken either from experimental studies or from ab initio calculations. In the present calculation, when possible, empirical potentials were used. The diatomic potentials were expressed analytically by the equation

$$U = U_0 + A\exp[-\alpha R] - C_6/R^6 - C_4/R^4 - C_1/R, \quad (44)$$

where R is the interatomic distance, U_0 is the energy of two separated atoms or ions, C_6 is the van der Waals coefficient, C_4 is the polarization coefficient [Eq.(13)], and $C_1 = e^2$ is the coefficient of the electrostatic attraction between two opposite charges. In the case of a pair of neutral atoms, only the repulsive and van der Waals terms remain since $C_1 = C_4 = 0$:

$$U = U_0 + A\exp[-\alpha R] - C_6/R^6 \quad (44')$$

The \bar{U} potential [Eq.(16)] of an ion-atom pair is also expressed by Eq. (44'), since $C_1 = 0$ and the polarization term is excluded. For a better fitting, some of the potentials can include two exponential functions or the $-C_8/R^8$ term. The exchange terms V are expressed by an exponential function

$$V = A\exp[-\alpha R]. \quad (45)$$

The parameters of Eqs. (44) and (45) are presented in Table 1. The sources of the potentials are given below.

A. Xe-Xe potential

The parameters of the expression (44') for the ground state Σ potential were obtained by the interpolation of the experimental data of the Xe-Xe interaction.⁴²

B. (Xe-Xe)⁺ potentials and exchange terms

The ab initio Xe_2^+ potential curves Σ_g , Σ_u , Π_u , Π_g without spin-orbit coupling⁴³ were used to calculate the Coulombic potentials U [Eqs.(35) and (37)] and the exchange terms V [Eqs. (36) and (38)]. These potentials are

preferable as they provide the ground-state potential well ($D_e = 1.08\text{eV}$, $R_e = 3.22 \text{ \AA}$) which is more reliable from the point of view of experimental results⁴⁴ than that obtained in Ref. 43 by including the spin-orbit coupling. The $(\text{Xe-Xe})^+$ diatomic terms, U and V, were interpolated by the analytical expressions, Eqs.(44) and (45).

C. Xe-Cl⁻ potential

The Xe-Cl⁻ Σ potential was obtained by interpolation of experimental data.⁴⁵ In order to obtain better fitting for short distances, the repulsive term is described by two exponential functions.

D. Xe-Cl potentials and coupling terms (Fig. 1)

In order to find the Xe-Cl Coulombic terms $U^{(00)}$ and $U^{(+-)}$, one needs the Xe-Cl potentials and the coupling term V [see the Eqs. (41)-(42)]. The potential curves of the ground state $1\Sigma(X)$ and the excited ionic state $2\Sigma(B)$ were obtained by the interpolation of experimental data.²⁶ The potential curve of the excited ionic $2\Pi(C)$ state is similar to the $2\Sigma(B)$ potential curve.²² The equilibrium point energy of 2Π was found experimentally to be almost the same as in the 2Σ state⁴⁶ or to be lower by only 0.016 eV ⁸ or 0.028 eV .⁴⁷ The 2Π equilibrium distance was found to be $R_e = 3.074 \text{ \AA}$,⁴⁷ which is larger by -0.06 \AA than that in the 2Σ state. The 2Π potential curve used in this paper has a minimum at the point $R_e = 3.074 \text{ \AA}$ and minimum energy value lower than the 2Σ state by 0.02 eV , which is a compromise of different experimental results.^{8,46,47} The polarization coefficient C_4 for the 2Π potential is assumed to be the same as for the 2Σ potential curve.

The potential curve of the first excited $1\Pi(A)$ state was found to have a minimum at the point $R_e = 4.1 \text{ \AA}$.⁴⁸ The 1Π potential curve used in the present paper fits this minimum point and the energy of transition from the

equilibrium point of the 2Π state to the repulsive 1Π state, which is equal to 3.59 eV (345 nm).⁴⁹

By using empirical Xe-Cl potentials we take into account indirectly the spin-orbit coupling, but only as a shift of the diatomic terms in the diagonal matrix elements. Any influence of the interaction of the Xe-Cl fragment with other atoms on the spin-orbit coupling is neglected. Besides, the consideration is restricted to states which correspond asymptotically to the Xe^+ and Cl atoms in their $P_{3/2}$ state.

As mentioned in the previous section, the coupling terms V_{Σ} and V_{Π} which are responsible for the mixing of the neutral and ionic states can be considered as semiempirical parameters. In the present calculation they were determined by fitting the XeCl transition moments. The ab initio transition moments were found as functions of interatomic distances.²² The empirical transition moments can be found at the equilibrium points of the ionic states by using the radiative lifetime τ_{kl} ²²,

$$\mu_{kl} = 77.9(E_k - E_l)^{-3/2} \sqrt{\tau_{kl}}, \quad E_k > E_l \quad (46)$$

where μ is expressed in Debyes (D), E in eV, and τ in nsec. The lifetimes of the B \rightarrow X and C \rightarrow A (Fig. 1) transitions are equal to 11.1 ns and 131 ns, respectively.⁴⁹ Substituting these values in the expression (46), as well as the energy differences (4.02 eV for B \rightarrow X and 3.59 eV for C \rightarrow A transition), one obtains $\mu_{\Sigma} = 2.9$ D for the B \rightarrow X transition moment and $\mu_{\Pi} = 1.0$ D for the C \rightarrow A transition moment. These values were fitted by the coupling terms V_{Σ} and V_{Π} presented as exponential functions [Eq.(45)]. The coefficients α of these exponential functions were determined by using the ab initio transition moments.²² The parameters of the coupling functions are given in

Table 1. The dependence of the B-X and C-A transition moments on the Xe-Cl distance is demonstrated in Fig. 2a.

The effect of the coupling on the Xe-Cl interaction is demonstrated in Fig. 2b by presenting the coupling energy, which is equal to the difference between the Coulombic potential $U_{\Sigma}^{(00)}$ and the ground-state potential 1Σ , and the contribution (in %) of the ionic Xe^+Cl^- configuration to the ground state. Both the coupling energy and the ionic configuration contribution decrease exponentially as the internuclear distance increases. At the point of the van der Waals equilibrium (3.23 Å), the coupling energy is 0.10 eV, almost equal to the energy of the van der Waals attraction (0.12 eV). The contribution of the ionic configuration at this point is 2.4%.

V. Results of calculations

The semiempirical DIIS calculations were performed for the following Xe_nCl systems: Xe_2Cl , Xe_3Cl and Xe_{12}Cl . The Xe_2Cl system has been studied both experimentally^{7,9,11,12} and theoretically.^{28,30} In the present calculation, which unlike another semiempirical calculation³⁰ takes into account directly the neutral-ionic coupling, the transition moments were obtained as well. The Xe_3Cl and Xe_{12}Cl systems are considered here in order to check the possibility of the formation of more stable ionic molecules than the Xe_2^+Cl^- molecule.

A. Xe_2Cl system

According to the results of the present calculation, the quasistable Xe_2^+Cl^- molecule which is formed in the first ionic state $4^2\Gamma$ has a configuration of an isosceles triangle, like in other Xe_2Cl calculations.^{28,30} The Xe-Cl distance in the Xe_2^+Cl^- molecule is 3.23 Å, or 0.16 Å larger than in the diatomic Xe^+Cl^- molecule. The Xe-Xe distance (3.25 Å) only slightly exceeds the interatomic distance in the Xe_2^+ molecule

(3.22 Å). The Xe_2Cl potential curves for a fixed Xe-Xe distance in the Xe_2^+Cl^- molecule (3.25 Å) are presented in Fig. 4a. The electronic states are denoted in the same way as in Ref. 30. Because of the neglect of the spin-orbit coupling, some of the states cannot be obtained in the present calculation. However, in contrast to Ref. 30, the calculation provides Π states with wave functions antisymmetric relatively to the molecule plane. The number of states is $3 \times 3 = 9$, including three largely neutral states and six largely ionic states. Three of these states ($2^2\Gamma_{\Pi}$, $4^2\Gamma_{\Pi}$, $6^2\Gamma_{\Pi}$) are the states of Π symmetry. The potentials of the antisymmetric states $4^2\Gamma_{\Pi}$ and $6^2\Gamma_{\Pi}$ almost coincide with the $5^2\Gamma$ and $6^2\Gamma$ potentials of the symmetrical states. The Xe_2Cl potential curves for fixed Xe-Cl distance of the Xe_2^+Cl^- molecule (3.23 Å) are presented in Fig. 4a. The antisymmetric states $2^2\Gamma_{\Pi}$, $4^2\Gamma_{\Pi}$ and $6^2\Gamma_{\Pi}$ and the highest state $8^2\Gamma$ are not shown in Fig. 4a. The transition moments for the pairs of states $1^2\Gamma-4^2\Gamma$, $2^2\Gamma-4^2\Gamma$ and $1^2\Gamma-5^2\Gamma$ are presented in Fig. 4b.

In the equilibrium configuration the energy of the Xe_2^+Cl^- molecule is 3.09 eV. The dominant transition from the $4^2\Gamma$ state is to the ground state $1^2\Gamma$ with the transition moment of 2.45 D, which is much higher than the ab initio transition moment of 0.96 D.²⁸ The transition moment to the state $2^2\Gamma$ has a sharp minimum in the Xe_2^+Cl^- equilibrium geometry and is very small (Fig. 4b). The calculated energy of the $4^2\Gamma-1^2\Gamma$ transition is 2.56 eV (485 nm), which is close to the experimental value of the center of the emission band (2.58 eV or 480 nm)¹¹ and to the ab initio value of 2.50 eV.²⁸ In the DIM calculation,³⁰ which was performed without taking into account directly the neutral-ionic coupling, the transition energy was found to be 2.74 eV. However, the difference between the transition energies in both calculations cannot be explained completely by the influence of the neutral-ionic

coupling, since in the DIM calculation³⁰ the ab initio diatomic potentials were used whereas the present study applies different, mainly empirical, diatomic potentials.

In order to check the influence of the coupling term V_{Σ} [see Eqs. (41)-(43)] on the results of calculation, the Xe_2^+Cl^- molecule was calculated also without the coupling term ($V_{\Sigma} = 0$). By putting $V_{\Sigma} = 0$ one does not neglect the neutral-ionic coupling since the empirical Xe-Cl diatomic potentials include it, but rather neglects only the dependence of the coupling energy on the system structure. The calculation shows that the neglect of the coupling term V_{Σ} decreases the energy of the excited ionic state by 0.07 eV, increases the energy of the ground state by 0.10 eV and, consequently, decreases the transition energy by 0.17 eV, which becomes equal to 2.38 eV. These energy changes demonstrate that the influence of the coupling term on the Xe_2Cl system does not change significantly the structure of the Xe_2Cl electronic states. Such relatively moderate influence of the coupling term on the Xe_2Cl can be explained by the fact that the mixing of neutral and ionic states in the Xe_2Cl system, as well as in other Xe_nCl systems, does not differ much from that of the XeCl diatomic system. For example, the electronic structure of the Xe_2^+Cl^- molecule includes 3.6% of the neutral electronic configuration, or 1.8% for one XeCl fragment. The corresponding value for the XeCl system with the same interatomic distance (3.23 Å) is 2.3% (Fig. 2b).

In addition to the "main" $4^2\Gamma \text{Xe}_2^+\text{Cl}^-$ molecule, there is also the quasistable collinear XeClXe molecule, which has Σ state symmetry and is 100% ionic. The Xe-Cl distances in this molecule are relatively short, 3.06 Å only. The Σ state molecule has 0.72 eV higher energy than the $4^2\Gamma \text{Xe}_2^+\text{Cl}^-$ molecule and is separated from $4^2\Gamma \text{Xe}_2^+\text{Cl}^-$ molecule by a high barrier. The

energy of transition to the ground state is 3.87 eV, and the transition moment is 3.9 D.

A quasistable molecule was found also in the ionic $4^2\Gamma_{\Pi}$ state. Like in the $4^2\Gamma$ state, the $4^2\Gamma_{\Pi}$ ionic molecule has a configuration of an isosceles triangle. The Xe-Cl distances are even shorter in this molecule (3.13 Å). However, the Xe-Cl-Xe angle is much larger (85°), and consequently the Xe-Xe distance (4.23 Å) significantly exceeds the Xe_2^+ equilibrium distance (3.22 Å). The energy of the $4^2\Gamma_{\Pi} \text{Xe}_2^+\text{Cl}^-$ molecule is 0.8 eV higher than that of the $4^2\Gamma \text{Xe}_2^+\text{Cl}^-$ molecule.

The ground state-potential energy surface is found to have two different van der Waals complexes with almost the same dissociation energy. One of these complexes, with the dissociation energy of 0.076 eV, has a symmetric XeClXe collinear configuration. The Xe-Cl distances are equal to 3.24 Å, which is 0.01 Å larger than in the XeCl diatomic complex. This slight increase in the Xe-Cl distance followed by an increase of the dissociation energy of 0.006 eV compared to the dissociation energy of two XeCl diatomic complexes can be explained only by the influence of the neutral-ionic coupling. The second van der Waals complex in the ground state, with the dissociation energy of 0.075 eV, has an asymmetric geometry. The Xe-Cl distances are 3.23 Å and 4.06 Å, and the Xe-Cl-Xe angle is equal to 74.4° . The energies of the excitation of the van der Waals symmetric (collinear) and asymmetric complexes to the ionic $4^2\Gamma$ state are equal to 3.96 eV and 3.94 eV, respectively, compared to the excitation energy of 4.05 eV for the case of the XeCl diatomic complex. The $1^2\Gamma \rightarrow 4^2\Gamma$ transition moment is large in the case of the collinear complex (3.32 D) but small (only 0.71 D) for the case of the asymmetric complex.

A van der Waals complex was found also in the first excited state ($2^2\Gamma$). It has a symmetrical geometry with the Xe-Cl distance of 3.99 Å and the Xe-Cl-Xe angle of 68.4°. The dissociation energy of this complex is positive, 0.055 eV.

The equilibrium configurations in the neutral states were presented above as van der Waals complexes. However, this term is not precise here as the attractive forces are provided also by the neutral-ionic coupling.

B. Xe₃Cl⁻ system

In order to check the possibility of the attachment of more Xe atoms to the Xe₂⁺Cl⁻ quasistable molecule, the Xe₃⁺Cl⁻ system was studied. The formation of the Xe₃⁺Cl⁻ molecules can be expected so far as the Xe₃⁺ molecules were found experimentally.⁵⁰ The most stable Xe₃⁺ configuration is a symmetric collinear one.⁵¹ According to the results of our calculation, the Xe₃⁺ arrangement in the Xe₃⁺Cl⁻ molecule is not collinear. In the most stable configuration, the Cl atom and two Xe atoms form an isosceles triangle with the Xe-Cl distance of 3.24 Å and the Xe-Xe distance of 3.25 Å, which almost coincide with the interatomic distances in the $4^2\Gamma$ Xe₂⁺Cl⁻ molecule. The third Xe atom is located in the plane perpendicular to the triangle plane at a distance of 3.52 Å from the Cl atom and 4.41 Å from the two other Xe atoms. The energy of dissociation of the Xe₃⁺Cl⁻ molecule to Xe₂⁺Cl⁻ and Xe is positive, 0.113 eV. It follows that the Xe₃⁺Cl⁻ can be formed, although the Xe₂⁺Cl⁻-Xe binding is weak and is determined mainly by the polarization interaction. The third Xe atom bears only 0.5% of the positive charge. The transition energy to the ground state is 2.48 eV. This value is not far from the Xe₂⁺Cl⁻ transition energy (2.56 eV), which makes it difficult to distinguish the Xe₃⁺Cl⁻ and Xe₂⁺Cl⁻ molecules by their emission.

A less stable Xe_3^+Cl^- molecule is formed in the symmetric configuration with the Xe-Cl distance of 3.33 Å and Xe-Xe distance of 3.58 Å. The dissociation energy in this configuration is 0.08 eV, and the transition energy is 2.62eV.

C. $\text{Xe}_{12}^+\text{Cl}^-$ system

According to the results of the calculation, the spherical symmetric $\text{Xe}_{12}^+\text{Cl}^-$ system is unstable. The stable configuration is formed by a Xe_2^+Cl^- molecule with slightly extended interatomic distances (3.29 Å for Xe-Cl and the same for Xe-Xe) and 10 weakly bounded Xe atoms. These 10 Xe atoms bear together only 4.3% of the positive charge. The energy of the dissociation of this system to Xe_2^+Cl^- and 10 Xe atoms is 0.88 eV. The energy of transition to the ground state is 2.49 eV. Considering the dissociation energies of the Xe_3Cl and Xe_{12}Cl systems, we have to note that they may be overestimated because of the neglect of three- and four-center interactions.⁵²

VI. Conclusion

1. The diatomic-in-ionic-systems (DIIS) method is designed for a semiempirical calculation of neutral polyatomic systems R_nX , which consist of atoms R of S symmetry and an atom X of P symmetry and which can form excited ionic R_n^+X^- states with positive ions R^+ of P symmetry and a negative ion X^- of S symmetry. Rare gas(R)-halogen(X) systems satisfy this definition and have been subject of consideration in the present paper. The DIIS method can be also used, with some modifications in the matrix element expressions, for calculations of other systems, such as hydrogen-containing rare gas-halides $\text{R}_n\text{H}_m\text{X}$, alkali metal-halides and rare gas cluster ions R_n^+ .

2. The DIIS method allows us to calculate the electronic structure of polyatomic systems by using diatomic potentials and transition moments as

initial data. In order to improve the results, it is possible also to use some known polyatomic values as semiempirical parameters. However the present calculation of polyatomic systems is based on diatomic data only. The calculation is limited to states without spin-orbit coupling. The influence of this coupling on the system energy is taken into account approximately and indirectly by using empirical diatomic potentials.

3. The DIIS method has been applied to the Xe_nCl systems, Xe_2Cl in particular. The ionic $Xe_2^+Cl^-$ in its most stable state ($4^2\Gamma$) was found to have an equilibrium geometry of an isosceles triangle, like in other calculations.^{28,30} In the equilibrium configuration the dominant transition is to the ground state, although in other configurations the radiative transition to the first excited state becomes possible as well. In addition to the $4^2\Gamma$ $Xe_2^+Cl^-$ molecule, there are other quasistable ionic molecules with higher energy, in particular, the collinear $Xe^{1/2+}Cl^-Xe^{1/2+}$ molecule. Any empirical evidence of the emission of these ionic molecules are unknown to us. In the ground state, two van der Waals complexes were found with the same dissociation energy but different geometries and different transition moments to the ionic states.

4. According to the results of the calculation, the ionic $Xe_2^+Cl^-$ molecule attracts more Xe atoms, but without sharing with them much of the Xe_2^+ positive charge. The ionic molecules with more than two Xe atoms, $Xe_3^+Cl^-$ in particular, can be formed in a dense gas. It is difficult to detect these molecules as their emission wavelength is similar to that of the $Xe_2^+Cl^-$ molecule.

5. The symmetric $Xe_{12}^+Cl^-$ clusters were found to be unstable, so that their formation in rare gas solids^{13,20} is unlikely.

Acknowledgments

This research was supported by the National Science Foundation under Grant CHE-8519053, the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United State Air Force, under Contract F49620-86-C-0009. The United State Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon. We are grateful to V. A. Apkarian for helpful discussions. I. L. thanks SUNY-Buffalo for its hospitality during a visit in 1986-87 when this work was performed.

References

1. S.K. Searles and G.A. Hart, Appl. Phys. Lett. 27, 243 (1975).
2. J.A. Mangano and J.H. Jacob, Appl. Phys. Lett. 27, 495 (1975).
3. J.J. Ewing and C.A. Brau, Appl. Phys. Lett. 27, 350 (1975); C.A. Brau and J.J. Ewing, J. Chem. Phys. 63, 4640 (1975).
4. J. Tellinghuisen, A.K. Hays, J.M. Hoffman and G.C. Tisone, J. Chem. Phys. 65, 4473 (1976).
5. V.S. Dubov, L.I. Gudzenko, L.V. Gurvich and S.I. Iakovlenko, Chem. Phys. Lett. 53, (1978).
6. D.C. Lorents, D.L. Huestis, M.V. McCusker, M.M. Nakano and R.M. Hill, J. Chem. Phys. 68, 4657 (1978).
7. F.K. Tittel, W.L. Wilson, R.E. Stickel, G. Marowski and W.E. Ernst, Appl. Phys. Lett. 36, 405 (1980)
8. J. Tellinghuisen and M.R. McKeeven, Chem. Phys. Lett. 72, 94 (1980).
9. V.S. Dubov, Y.E. Lapsker, A.N. Samoilova and L.V. Gurvich, Chem. Phys. Lett. 83, 518 (1981).
10. H.C. Brashears, Jr., D.W. Setser and Y.-C. Yu, J. Chem. Phys. 74, 10 (1981).
11. A.W. McCown and J.C. Eden, J. Chem. Phys. 81, 2933 (1984).
12. A.W. McCown, M.N. Ediger, D.B. Geohegan and J.G. Eden, J. Chem. Phys. 82, 4862 (1985).
13. M.E. Fajardo and V.A. Apkarian, J. Chem. Phys. 85, 5660 (1986)
14. M.E. Fajardo and V.A. Apkarian, J. Chem. Phys. (in press)
15. J. Tellinghuisen, J.M. Hoffman, G.C. Tisone and A.K. Hays, J. Chem. Phys. 64, 2484 (1976).
16. J.H. Waters and H.B. Gray, J. Am. Chem. Soc. 85, 825 (1963).
17. B.E. Wilcomb and R. Burnham, J. Chem. Phys. 74, 6784 (1981).
18. G.T. Fraser and A.S. Pine, J. Chem. Phys. 85, 2502 (1986).
19. N. Goldstein, T.L. Brack and G.H. Atkinson, J. Chem. Phys. 85, 2684 (1966).
20. I. Last and T.F. George, J. Chem. Phys. 86, (1987)
21. P.J. Hay and T.H. Dunning, Jr., J. Chem. Phys. 66, 1306 (1977).
22. P.J. Hay and T.H. Dunning, Jr., J. Chem. 69, 2209 (1978).

23. K. Balasubramanian, J.J. Kaufman, P.C. Hariharan and W.S. Koski, *Chem. Phys. Lett.* 129, 165 (1986).
24. Y.S. Kim and R.G. Gordon, *J. Chem. Phys.* 61, 1 (1974).
25. C. Vieugd, R.W.W. van Resandt and J. Los, *Chem. Phys. Lett.* 65, 93 (1979).
26. A. Sur, A.K. Hui and J. Tellinghuisen, *J. Mol. Spectrosc.* 74, 465 (1979).
27. W.R. Wadt and P.J. Hay, *J. Chem. Phys.* 68, 3850 (1978).
28. W.J. Stevens and M. Krauss, *Appl. Phys. Lett.* 41, 301 (1982).
29. D.L. Huestis and N.E. Schlotter, *J. Chem. Phys.* 69, 3100 (1978).
30. D.L. Huestis, G. Marowsky and F.K. Tittel, in Excimer Lasers - 1983, ed. by C.K. Rhodes, H. Egger and H. Pummer (American Institute of Physics, New York, 1983), pp. 238-248.
31. J.C. Tully, *Adv. Chem. Phys.* 42, 63 (1980).
32. J.J. Duggan and R. Grice, *J. Chem. Phys.* 78, 3842 (1983).
33. A.D. Isaacson and J.T. Muckerman, *J. Chem. Phys.* 73, 1729 (1980).
34. P.J. Kuntz and A.C. Roach, *J. Chem. Phys.* 74, 3420 (1981).
35. B.E. Douglas and D.H. McCanien, Concepts and Models of Inorganic Chemistry (Blasdel Publishing Company, New York, 1965), p. 97.
36. M. Mandick, P. Beeken and G. Flynn, *J. Chem. Phys.* 77, 702 (1982).
37. R. McWeeny, *Proc. Roy. Soc. A* 253, 242 (1959); R. McWeeny and B.T. Sutcliffe, Methods of Molecular Quantum Mechanics (Academic Press, New York, 1969).
38. E. Kapuy, *Acta Phys. Acad. Sci. Hung.* 11, 97 (1960).
39. I.Ts. Lyast, *J. Struct. Chem.* 16, 355 (1975).
40. I. Last, *Chem. Phys.* 55, 237 (1981).
41. N. Gresh, P. Claverie and A. Pullman, *Int. J. Quantum. Chem.* 29, 101 (1986).
42. J.S. Cohen and R.T. Pack, *J. Chem. Phys.* 61, 2372 (1974).
43. W.R. Wadt, *J. Chem. Phys.* 68, 402 (1978).
44. C.E. Melton and W.H. Hamill, *J. Chem. Phys.* 41, 1469 (1964); P.M. Dehmer, S.T. Pratt and J.L. Dehmer, *J. Chem. Phys.* 85, (1986); K.P. Killeen and J.G. Eden, *J. Chem. Phys.* 84, 6048 (1986).

45. Ch. C. Kirkpatrick and L.A. Viehland, Chem. Phys. 98, 221 (1985).
46. J. Bokor and C.K. Rhodes, J. Chem. Phys. 73, 2626 (1980).
47. Y.C. Yu, D.W. Setser and H. Horiguchi, J. Phys. Chem. 87, 2199 (1983).
48. C.H. Becker, J.J. Valentini, P. Casavecchia, S.J. Sibener and Y.T. Lee, Chem. Phys. Lett. 61, 1 (1979).
49. G. Inoue, J.K. Ku and D.W. Setser, J. Chem. Phys. 80, 6006 (1984).
50. H. Helm, Phys. Rev. A 14, 680 (1976).
51. U. Umehara, Phys. Rev. B 33, 4237 (1986).
52. I. Last and M. Baer, J. Chem. Phys. 75, 288 (1981).

Table 1

Parameters of diatomic terms. For the designation of the parameters see Eqs. (44) and (45). R_e and D_e are the equilibrium distance and the dissociation energy, respectively.

diatomic		U_0	α	A	C_6	C_4	C_1	R_e	D_e
fragment	term	eV	\AA^{-1}	eV	eV \AA^6	eV \AA^4	eV \AA	\AA	eV
Xe-Xe	Σ	0	3.17	24170	320	0	0	4.47	0.023
$\text{Xe}^+ \text{-Xe}$	U_Σ	0	2.95	14520	320	29.14	0	3.22*	1.08*
$(\text{Xe-Xe})^+$	V_Σ	-	1.45	-172	-	-	-	-	-
$\text{Xe}^+ \text{-Xe}$	U_Π	0	3.00	18625	320	29.14	0	4.00*	0.16*
$(\text{Xe-Xe})^+$	V_Π	0	1.90	-168	-	-	-	-	-
XeCl^-	Σ	0	3.85 8.0**	31670 $4 \cdot 10^8$	60	29.14	0	3.30	0.195
Xe-Cl	1Σ	0	3.10	1167	99.2	0	0	3.23	0.035
$\text{Xe}^+ \text{-Cl}^-$	2Σ	8.52	2.512	2033	0	66	14.4	3.01	453
Xe-Cl	1Π	0	3.90	86000	.2 310***	0	0	4.10	0.015
$\text{Xe}^+ \text{-Cl}^-$	2Π	8.52	2.847	5520	0	66	14.4	3.074	3.97
Xe-Cl	V_Σ	-	1.20	-30	-	-	-	-	-
Xe-Cl	V_Π	-	1.40	-18	-	-	-	-	-

* R_e and D_e for Xe_2^+ potential curves connected with U and V by Eqs. (35)-(38).

** Parameters of the second exponent of the XeCl^- potential.

*** Coefficient C_8 of the C_8/R^8 term of the 1Π potential.

Figure Captions

Fig. 1. XeCl potential curves (Table 1).

Fig. 2. The effects of coupling between ionic and neutral electronic configurations in the XeCl system. The vertical dashes stand for the 2Π (C) equilibrium distance.

a) Stabilization energy of the coupling (right ordinate) and the contribution of ionic configuration (left ordinate) in the ground 1Σ (X) state.

b) $1\Sigma-2\Sigma$ (X-C) and $1\Pi-2\Pi$ (A-C) transition moments.

Fig. 3. Xe_2Cl potential energy surfaces for a geometry (shown in the figure) with a fixed Xe-Xe distance equal to that of the $Xe_2^+Cl^-$ ($4^2\Gamma$ state) molecule. The vertical dashes stand for the equilibrium $Xe_2^+Cl^-$ configuration.

Fig. 4. Xe_2Cl system in a geometry (shown in the figure) with a fixed Xe-Cl distance equal to that of the $Xe_2^+Cl^-$ ($4^2\Gamma$ state) molecule. The vertical dashes stand for the equilibrium $Xe_2^+Cl^-$ configuration.

a) potential energy surfaces for five $^2\Gamma$ states.

b) $1^2\Gamma-4^2\Gamma$, $1^2\Gamma-5^2\Gamma$ and $2^2\Gamma-4^2\Gamma$ transition moments.

Fig. 1




Fig. 2(a)






Fig. 2(b)





(a)





Fig. 4 (b)



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburg
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 41106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853

END

5-87

DTIC